

Set Name **Query**
side by side

Hit Count **Set Name**
result set

DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

<u>L17</u>	L16 with l15 with l14 with l6	27	<u>L17</u>
<u>L16</u>	l1 or l4	1417861	<u>L16</u>
<u>L15</u>	AQUEOUS	875317	<u>L15</u>
<u>L14</u>	alcohol or PVA	653344	<u>L14</u>
<u>L13</u>	L12 with l6	3	<u>L13</u>
<u>L12</u>	l4 with l3	53	<u>L12</u>
<u>L11</u>	L10 with l5	16	<u>L11</u>
<u>L10</u>	L8 with l3	3445	<u>L10</u>
<u>L9</u>	L8 and l3	8041	<u>L9</u>
<u>L8</u>	L7 or l1	1452645	<u>L8</u>
<u>L7</u>	peg	68146	<u>L7</u>
<u>L6</u>	dna or plasmid or nucleic or polynucleotide	181920	<u>L6</u>
<u>L5</u>	dna or plasmid or nucleic	175535	<u>L5</u>
<u>L4</u>	microparticle or microsphere	33241	<u>L4</u>
<u>L3</u>	negatively charged lipid or DPSE or DOPE	40676	<u>L3</u>
<u>L2</u>	s negatively charged lipid or DPSE or DOPE	40411	<u>L2</u>
<u>L1</u>	s peg or polymer	1403779	<u>L1</u>

END OF SEARCH HISTORY

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)[Cases](#)**Search Results -**

Terms	Documents
L16 with I15 with I14 with I6	27

Database:
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Search:[Refine Search](#)[Recall Text](#)[Clear](#)**Search History**

DATE: Tuesday, January 14, 2003 [Printable Copy](#) [Create Case](#)

WEST

Generate Collection

Print

L17: Entry 2 of 27

File: PGPB

Jun 27, 2002

DOCUMENT-IDENTIFIER: US 20020082237 A1

TITLE: Cationic polymers and lipids for the delivery of nucleic acids


Summary of Invention Paragraph (29):

[0024] Another embodiment of the present invention includes a novel process for making polynucleotide delivery vehicles comprising the steps of complexing the polynucleotide and cationic polymer and/or cationic lipid in buffer that maintains DNA as a B-form helix (e.g., an aqueous alcohol solution), and removing the buffer by evaporation. After reconstitution of the dried polynucleotide-cationic lipid/cationic polymer complex with aqueous solution, stable polynucleotide delivery vehicles are produce.

(FILE 'MEDLINE, CANCERLIT, EMBASE, BIOSIS, BIOTECHDS' ENTERED AT 14:03:35
ON 14 JAN 2003)

DEL HIS

L23	873716	S POLYME? OR MICROPARTICLE OR MICROSPHERE
L24	536386	S LIPID OR AMPHIPHILE
L25	9979	S L24 AND L23
L26	2369734	S DNA OR NUCLEIC OR PLASMID
L27	2961	S L26 AND L25
L28	30437	S PEG
L29	33	S L28 AND L27
L30	14	DUP REM L29 (19 DUPLICATES REMOVED)
L31	1987	S PLG
L32	5	S L31 AND L27
L33	2	DUP REM L32 (3 DUPLICATES REMOVED)
L34	4660255	S MICRO?
L35	758	S L34 AND L27
L36	552	DUP REM L35 (206 DUPLICATES REMOVED)
L37	35164	S ANIONIC OR NEGATIVELY CHARGED LIPID
L38	3822	S L37 AND L24
L39	413	S DSPE
L40	10	S L39 AND L38
L41	5	DUP REM L40 (5 DUPLICATES REMOVED)
L42	53	S L39 AND L25
L43	27	DUP REM L42 (26 DUPLICATES REMOVED)
L44	40	SS L27 AND (L37 OR L39)
L45	23	DUP REM L44 (17 DUPLICATES REMOVED)
L46	2932	S MICROPARTICLE
L47	16	S L46 AND L27
L48	11	DUP REM L47 (5 DUPLICATES REMOVED)

**PALM INTRANET**Day : Tuesday
Date: 1/14/2003
Time: 15:46:41

Continuity Information for 09/909460

Parent Data

09909460is a continuation of 09321346Which is a continuation in part of 09266463Which is a continuation in part of 09003253Which Claims Priority from Provisional Application 60035983Which is a continuation in part of PCT/US98/01499 International Filing Date: 01/22/1998

Child Data

No Child Data

Appln Info

Contents

Petition Info

Atty/Agent Info

**Continuity
Data**

Foreign Data

Inventors

Search Another: Application# or Patent# PCT / / or PG PUBS # Attorney Docket #

To go back use Back button on your browser toolbar.

Back to [PALM](#) | [ASSIGNMENT](#) | [OASIS](#) | [Home page](#)

L30 ANSWER 12 OF 14 MEDLINE DUPLICATE 8
 AN 1999337493 MEDLINE
 DN 99337493 PubMed ID: 10407066
 TI Stabilized **plasmid-lipid** particles: factors influencing **plasmid** entrapment and transfection properties.
 AU Mok K W; Lam A M; Cullis P R
 CS Department of Biochemistry and Molecular Biology, University of British Columbia, 2146 Health Sciences Mall, Vancouver, B.C. V6T 1Z3, Canada.
 SO BIOCHIMICA ET BIOPHYSICA ACTA, (1999 Jul 15) 1419 (2) 137-50.
 Journal code: 0217513. ISSN: 0006-3002.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199909
 ED Entered STN: 19990921
 Last Updated on STN: 19990921
 Entered Medline: 19990903
 AB Previous work has shown that **plasmid DNA** can be encapsulated in small 'stabilized **plasmid-lipid** particles' (SPLP) composed of 1, 2-dioleoyl-3-phosphatidylethanolamine (DOPE), the cationic **lipid** N, N-dioleoyl-N,N-dimethylammonium chloride (DODAC) and poly(ethylene glycol) (**PEG**) conjugated ceramides (**PEG-Cer**), employing a detergent dialysis procedure. These SPLP have potential as vectors for in vivo gene therapy. This study is aimed at characterizing the influence of the cationic **lipid** and **PEG-Cer** species on SPLP formation and in vitro transfection properties. It is shown that the transfection potency of SPLP is sensitive to the cationic **lipid** species employed, the size of the **PEG polymer** incorporated in the **PEG-ceramide** and the length of the acyl chain contained in the ceramide anchor. With regard to the influence of cationic **lipid**, the transfection levels achieved were highest for SPLP containing N-[2, 3-(dioleoyloxy)propyl]-N,N-dimethyl-N-cyanomethylammonium chloride (DODMA-AN) and lowest for SPLP containing 3-beta-[N-(N', N'-dimethylaminoethyl)carbamoyl]-cholesterol (DC-CHOL), according to the series DODMA-AN>N-[2,3-(dioleoyloxy)propyl]-N,N, N-trimethylammonium chloride (DOTMA)>DODAC>N,N-distearyl-N, N-dimethylammonium chloride (DSDAC)>DC-CHOL. Incorporation of short (**PEG**(750)) **PEG polymers** in the **PEG-ceramide** components resulted in modest improvements in transfection levels over **PEG**(2000) and **PEG**(5000) **polymers**, however variation of the length of the acyl chain contained in the hydrophobic ceramide anchor from octanoyl (**PEG-CerC**(8)) to myristoyl (**PEG-CerC**(14)) to arachidoyl (**PEG-CerC**(20)) had the most dramatic effects. Transfection levels achieved for SPLP containing **PEG-CerC**(8) were substantially larger than observed for SPLP containing **PEG-CerC**(14) or **PEG-CerC**(20), consistent with a requirement for the **PEG-ceramide** to dissociate from the SPLP surface for maximum transfection potency. It is also shown that the ability of SPLP to be accumulated into cells is a dominant factor influencing transfection potency, and that the transfection potency of SPLP that are accumulated is at least equivalent to that of cationic **lipid-plasmid DNA** complexes.

L30 ANSWER 11 OF 14 MEDLINE DUPLICATE 7
 AN 1999363970 MEDLINE
 DN 99363970 PubMed ID: 10435112
 TI Stabilized **plasmid-lipid** particles: construction and characterization.
 AU Wheeler J J; Palmer L; Ossanolou M; MacLachlan I; Graham R W; Zhang Y P; Hope M J; Scherrer P; Cullis P R
 CS Inex Pharmaceuticals Corporation, Burnaby, BC, Canada.
 SO GENE THERAPY, (1999 Feb) 6 (2) 271-81.
 Journal code: 9421525. ISSN: 0969-7128.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199908
 ED Entered STN: 19990827
 Last Updated on STN: 19990827
 Entered Medline: 19990817
 AB A detergent dialysis procedure is described which allows encapsulation of **plasmid DNA** within a **lipid** envelope, where the resulting particle is stabilized in aqueous media by the presence of a poly(ethyleneglycol) (**PEG**) coating. These 'stabilized **plasmid-lipid** particles' (SPLP) exhibit an average size of 70 nm in diameter, contain one **plasmid** per particle and fully protect the encapsulated **plasmid** from digestion by serum nucleases and E. coli DNase I. Encapsulation is a sensitive function of cationic **lipid** content, with maximum entrapment observed at dioleoyldimethylammonium chloride (DODAC) contents of 5 to 10 mol%. The formulation process results in **plasmid**-trapping efficiencies of up to 70% and permits inclusion of 'fusigenic' lipids such as dioleoylphosphatidylethanolamine (DOPE). The in vitro transfection capabilities of SPLP are demonstrated to be strongly dependent on the length of the acyl chain contained in the ceramide group used to anchor the **PEG polymer** to the surface of the SPLP. Shorter acyl chain lengths result in a **PEG** coating which can dissociate from the SPLP surface, transforming the SPLP from a stable particle to a transfection-competent entity. It is suggested that SPLP may have utility as systemic gene delivery systems for gene therapy protocols.

L30 ANSWER 2 OF 14 MEDLINE DUPLICATE 1
 AN 2002200333 MEDLINE
 DN 21848587 PubMed ID: 11858858
 TI Protective immune responses elicited in mice by immunization with
 formulations of poly(lactide-co-glycolide) microparticles.
 AU McKeever U; Barman S; Hao T; Chambers P; Song S; Lunsford L; Hsu Y-Y; Roy
 K; Hedley M L
 CS Zycos Inc., 44 Hartwell Avenue, Lexington, MA 02421, USA.
 SO VACCINE, (2002 Feb 22) 20 (11-12) 1524-31.
 Journal code: 8406899. ISSN: 0264-410X.
 CY England: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200209
 ED Entered STN: 20020406
 Last Updated on STN: 20020928
 Entered Medline: 20020927
 AB Parenteral administration of **microparticle** encapsulated
DNA elicits immune responses to the encoded antigens. Experiments
 were performed to test whether the addition of certain lipophilic agents
 to such formulations enhanced the activity of a beta-galactosidase
 (beta-gal) **DNA** vaccine. Addition of either taurocholic acid (TA)
 or monomethoxy polyethylene-glycol-distearoylphosphatidylethanolamine (
PEG-DSPE) increased the efficiency of **DNA** encapsulation.
 Immunization of mice with encapsulated **DNA** formulations
 containing either compound significantly increased the number of antibody
 positive responders over that achieved with non-**lipid** containing
 particles. Moreover, responding animals demonstrated trends towards higher
 antibody titers and increased T cell responses. Tumor protection against
 the CT26.CL25 tumor cell line was demonstrated with **lipid** and
 non-**lipid** containing formulations. These results are the first
 demonstration of protection obtained by parenteral administration of PLG
 encapsulated **DNA** vaccines.

L33 ANSWER 1 OF 2 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI
 AN 2002-11585 BIOTECHDS
 TI Novel **nucleic** acid delivery system useful for preparation of composition for delivering **nucleic** acid to subject and for treating/preventing cancer, comprises **DNA** encapsulated in biodegradable **polymeric** microspheres;
 recombinant vector-mediated gene transfer and expression in host cell for use in recombinant vaccine and **nucleic** acid vaccine preparation and cancer prevention, therapy and gene therapy
 AU JOHNSON M E; MOSSMAN S; CECIL T; EVANS L
 PA CORIXA CORP
 PI WO 2002003961 17 Jan 2002
 AI WO 2000-US21780 7 Jul 2000
 PRAI US 2000-216604 7 Jul 2000
 DT Patent
 LA English
 OS WPI: 2002-257248 [30]
 AB DERWENT ABSTRACT:
 NOVELTY - A **nucleic** acid delivery system (I) comprising deoxyribonucleic acid (**DNA**) encapsulated in biodegradable **polymeric** microspheres, where at least 50% of the **DNA** comprises supercoiled **DNA**, and where at least 50% of the **DNA** is released from the microspheres after 7 days at about 37 degrees C, is new.
 DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the following: (1) encapsulating (M) **nucleic** acid molecules in microspheres by dissolving a **polymer** in a solvent to form a **polymer** solution, adding an aqueous solution containing **nucleic** acid molecules to the **polymer** solution to form a primary emulsion, homogenizing the primary emulsion, mixing the primary emulsion with a process medium comprising a stabilizer to form a secondary emulsion, and extracting the solvent from the secondary emulsion to form microspheres encapsulating **nucleic** acid molecules; (2) a pharmaceutical composition (II) comprising **nucleic** acid molecules encapsulated in microspheres produced by (M); and (3) use of an aminoalkyl glucosaminide 4-phosphate (AGP) (III) for the preparation of an adjuvant for enhancing the immunostimulatory efficacy of microspheres encapsulating **nucleic** acid molecules.
 WIDER DISCLOSURE - Also disclosed is an adjuvant for modulating the immunostimulatory efficacy of microspheres encapsulating **nucleic** acid molecules comprising aminoalkyl glucosaminide 4-phosphate (AGP).
 BIOTECHNOLOGY - Preferred System: In (I), the microspheres have an encapsulation efficiency of at least about 40%. At least about 70% of the **DNA** is released from the microspheres after 7 days at about 37 degrees C. At least about 90% of the microspheres are about 1-10 microm in diameter. The microspheres comprise poly(lacto-co-glycolide) (**PLG**). (I) further comprises an adjuvant comprising AGP. The **DNA** encodes an antigen such as her2/neu associated with cancer, preferably breast cancer or an antigen such as TbH9 associated with infectious disease, preferably tuberculosis. Preferred Method: In (M), the **polymer** comprises **PLG** including ester end groups or carboxylic acid end groups. The **PLG** has a molecular weight of from about 8-65 kDa. The **nucleic** acid molecules are maintained at about 2-35 degrees C, preferably 4-25 degrees C prior to the extraction. The solvent comprises dichloromethane, chloroform, or ethylacetate. The **polymer** solution further comprises a cationic **lipid**, and an adjuvant comprising MPL. The stabilizer comprises carboxymethylcellulose (CMC), polyvinyl alcohol (PVA), or a mixture of CMC and PVA, and a cationic **lipid**. The stabilizer comprises from about 1-5% of the process medium. The solvent comprises an internal water volume of from about 0.001-0.5%. The aqueous solution comprises an

ethanol content of from about 0-75% (v/v). The **nucleic acid** molecule comprises **DNA**. The aqueous solution comprises about 0.2-12 mg/ml **DNA** comprising a **plasmid** of about 3-9 kb. The aqueous solution further comprises an adjuvant such as QS21, and a stabilizer comprising bovine serum albumin. At least 50% of the **DNA** retains a supercoiled formation through the extraction step. The encapsulation efficiency is at least about 40%. The microspheres release at least about 50% of the **nucleic acid** molecules within about 7 days, preferably 4 days. Preferred Composition: (II) further comprises an adjuvant such as AGP. Preferred Adjuvant: (III) comprises an aqueous formulation. (III) is preferably 517, 527, 547, 557 or 568. (III) is administered simultaneously with the microspheres, or before or after administration of the microspheres.

ACTIVITY - Cytostatic; tuberculostatic.

MECHANISM OF ACTION - Enhancer of immunostimulatory efficacy of microspheres encapsulating **nucleic acid** molecules (claimed); vaccine; gene therapy. Immune responses elicited in monkeys by encapsulated **DNA** was tested: The immune responses elicited in rhesus macaques following three immunizations, at monthly intervals, with either naked TbH9-VR1012 **DNA** or TbH9-VR1012 **DNA** encapsulated in microspheres were tested. Naked **DNA** consisted of 3.3 mg **plasmid** + 40 microg RC527-AF, immunized by intradermal and intramuscular routes. Microspheres **DNA** consisted of 3 mg **plasmid** + 50 microg RC 568-Af delivered intramuscularly. There were four animals in each group. The results, demonstrated that the **microsphere**-encapsulated **DNA** elicited stronger immune responses than were observed with naked **DNA**.

USE - (I) is useful for the preparation of a composition for delivering a **nucleic acid** molecule to a subject, for eliciting an immune response to an antigen in a subject, for treating or preventing a cancer associated with her2/neu antigen or tuberculosis in a subject. (III) is useful for the preparation of an adjuvant for enhancing the immunostimulatory efficacy of microspheres encapsulating **nucleic acid** molecules (claimed). (I) is useful for delivery of vaccines, preferably **DNA** vaccines.

ADMINISTRATION - (II) is administered by parenteral (e.g., intravenous, subcutaneous, intramuscular), buccal, sublingual, rectal, oral, nasal, topical (e.g., transdermal, ophthalmic), vaginal, pulmonary, intraarterial, intraperitoneal, intraocular, or intranasal route or directly into a specific tissue. No dosage details are given.

ADVANTAGE - (I) offers, in one system, a combination of high encapsulation efficiency, rapid release kinetics and preservation of **DNA** in supercoiled form.

EXAMPLE - The formulation of a **DNA** poly(lacto-co-glycolide) (**PLG**) microspheres with desirable in vitro characteristics was as follows. Specifically, 1-10 microm diameter microspheres which were able to release their **DNA** contents over the course of a week were prepared using a process that resulted in a high encapsulation efficiency (60-80%) and high rate of retention of the **DNA** supercoiled state (70%). **PLG** microspheres containing **DNA** encoding antigenic proteins were prepared using variations on the double emulsion technique (J.H. Eldridge et al. Mol Immunol, 28:287-294, 1991; S. Cohen et al. Pharm Res, 8:713-720, 1991)). Specifically, **plasmid DNA** in Tris-ethylenediaminetetraacetic acid (EDTA) buffer, 0.38 ml ethanol were combined and brought up to a volume of 5.1 ml using Tris-EDTA buffer. This was the internal (water) phase. 1200 mg of **PLG** **polymer** was dissolved in 13.9 ml of dichloromethane (DCM) and put on ice. The internal aqueous phase was added to the **PLG** solution and mixed in a 30 ml syringe while still on ice using a Polytron tissue homogenizer for 20 seconds to form the primary emulsion

(water-in-oil). The secondary emulsion was prepared by adding the primary emulsion to a beaker containing 280 ml of 1.4% carboxymethylcellulose), or process medium, on ice, and mixing. The secondary emulsion was diluted with miliQ water, and mixed in order to extract dichloromethane from, and to harden, the microspheres. The resulting microspheres were washed and centrifuged. After washing, mannitol was added to the concentrated microspheres, which were frozen and lyophilized. Lyophilized microspheres were then assayed for their size distribution, **DNA** content, release kinetics, and the supercoiled content of the encapsulated **DNA**. Two plasmids were used in this study, one encoding a tuberculosis antigen, TbH9, and the other encoding the breast cancer antigen, Her-2/neu. Mice were immunized with **DNA** microspheres dispersed in aqueous buffer. The combination of microspheres with selected aminoalkyl glucosaminide 4-phosphate (AGP) was investigated by using a sub-optimal immunization schedule, a single 10 microg dose of encapsulated **DNA** dispersed in phosphate buffer saline (PBS) along with 10 microg of adjuvant. Lastly, the effect of the resuspension buffer was examined by administering to mice a single 10 microg dose of encapsulated **DNA** dispersed in either PBS or sodium chloride free phosphate buffer (PB). The process resulted in microspheres that were small (about 1-10 microm in diameter), with rapid release kinetics, high encapsulation efficiency (40-80%), and good retention of supercoiled **DNA**. More than 33% of the **microsphere** contents were released after 48 hours in vitro at 37 degrees C, more than 50% were released after 4 days, and more than 70% after 7 days. The ratio of supercoiled-to-nicked **DNA** for the **plasmid** extracted from the microspheres was more than 50% of the ratio of the input **DNA**. (60 pages)

L41 ANSWER 5 OF 5 MEDLINE DUPLICATE 3
 AN 2000419156 MEDLINE
 DN 20302306 PubMed ID: 10845683
 TI Novel therapeutic nano-particles (lipocores): trapping poorly water soluble compounds.
 AU Perkins W R; Ahmad I; Li X; Hirsh D J; Masters G R; Fecko C J; Lee J; Ali S; Nguyen J; Schupsky J; Herbert C; Janoff A S; Mayhew E
 CS The Liposome Company, Inc., Princeton, NJ 08540, USA.. wperkins@lipo.com
 SO INTERNATIONAL JOURNAL OF PHARMACEUTICS, (2000 Apr 25) 200 (1) 27-39.
 Journal code: 7804127. ISSN: 0378-5173.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200009
 ED Entered STN: 20000915
 Last Updated on STN: 20000915
 Entered Medline: 20000901
 AB The development of stable spherical **lipid**-coated drug particles that are termed 'lipocores' is reported here. Unlike conventional **lipid**-based particles (i.e. liposomes, emulsions, micelles), these particles are comprised solely of a core of a poorly water soluble drug surrounded by polyethyleneglycol conjugated **lipid** (PEG-**lipid**) and are formed via a 'kinetic' trapping process. These lipocore particles were made with the acyl chain of 16 carbon length (C16) acyl-chain derivatives of paclitaxel or vinblastine and with the polyene antifungal hamycin. Formation of the particles occurred regardless of the type of PEG-phospholipid used (i.e. acyl chain length, chain saturation, and polymer length) and could also be formed with the **negatively charged lipid** N-glutaryl-dioleoyl-phosphatidylethanolamine (DOPE-GA). Images from both freeze-fracture electron microscopy and electron cryo-microscopy revealed solid spherical structures with no internal lamellae for the PEG-**lipid** particles made with the C16 derivatives of paclitaxel (BrC16-T) or vinblastine (C16-Vin). From a solute distribution study of lipocores made with BrC16-T and distearoyl-phosphatidylethanolamine-PEG2000 (**DSPE**-PEG2000), the particles were found to have no measurable aqueous captured volume. Fluorescence anisotropy and order parameter measurements revealed the core material of these particles to be highly immobilized. The mole ratio of BrC16-T:**lipid** in the lipocores was typically > 90: < 10 and as high as 98:2, and the refrigerated lipocores were stable for several months. BrC16-T/**DSPE**-PEG2000 lipocores of 50-100 nm particle size were far less toxic than paclitaxel (Taxol) after intraperitoneally (i.p.) or intravenously (i.v.) administration in mice and were active against i.p. and subcutaneously (s.c.) planted human (OvCar3) ovarian carcinoma grown in SCID mice. It is believed the high drug:**lipid** ratio, the stability, and therapeutic efficacy of these novel particles make them a paradigm for delivery of poorly water soluble drugs and/or their hydrophobic derivatives.

L41 ANSWER 2 OF 5 MEDLINE
 AN 2003013680 IN-PROCESS
 DN 22407987 PubMed ID: 12519628
 TI THE TRANSFECTION OF JURKAT T-LEUKEMIC CELLS BY USE OF pH-SENSITIVE IMMUNOLIPOSOMES.
 AU Turner Christopher; Weir Neil; Catterall Catherine; Baker Terry S; Carrington Bruce; Jones Malcolm N
 CS Liposome Research Unit, University of British Columbia, Vancouver, BC, V6T 123, Canada.
 SO J Liposome Res, (2002 Nov) 12 (4) 311-34.
 Journal code: 9001952. ISSN: 0898-2104.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS IN-PROCESS; NONINDEXED; Priority Journals
 ED Entered STN: 20030110
 Last Updated on STN: 20030110
 AB A gene transfer vector has been developed utilising **anionic** liposomes as a carrier of plasmid DNA (pEGlacZ, 7.6 kb) to transfect CD3+ T lymphocytes (Jurkat cells). The plasmid DNA that contained the Escherichia coli beta-galactosidase reporter gene was condensed using poly-l-lysine of molecular mass 20,700 (PLK99) to form a polyplex which was interacted with several **anionic** liposome formulations to form lipopolyplexes. The liposome formulations were based on dioleoylphosphatidylethanolamine (DOPE) in combination with cholesterol and dioleoylphosphatidylcholine (DOPC) and oleic acid, or dimyristoylphosphatidylethanolamine (DMPE). For targeting to the Jurkat cells distearoylphosphatidylethanolamine (**DSPE**) linked to poly (ethylene glycol) molecular mass 2000 and coupled to anti-CD3 antibody was incorporated. The polyplexes and lipopolyplexes were characterised in terms of size, zeta potential, agarose gel electrophoresis and electron microscopy and the permeability of the lipopolyplexes to liposome-encapsulated glucose was determined. The polyplexes consisted of a mixed population of rod-like structures (53-160 nm long and 23-31 nm diameter) and spheres (18-30 nm diameter). The lipopolyplexes retained a permeability barrier although were more permeable to glucose than their component liposomes. The poly-l-lysine condensing agent was still susceptible to pronase digestion suggesting that the polyplex was associated with the outer surface of the liposome. The lipopolyplexes with **lipid** composition DOPE/cholesterol/OA/**DSPE**-PEG2000 anti-CD3+ PLK99-plasmid DNA had significant gene transfer activity, as monitored by beta-galactosidase expression, that depended on the charge ratio of the component polyplex and the **lipid**/DNA weight ratio. The anti-CD3 antibody, the liposomal **lipid** and pH sensitivity were essential for transfection activity.

L43 ANSWER 27 OF 27 EMBASE COPYRIGHT 2003 ELSEVIER SCI. B.V.
 AN 93131099 EMBASE
 DN 1993131099
 TI Rationale for the control of the biological properties of **lipid** membranes in vitro and in vivo.
 AU Cevc G.; Blume G.
 CS Medizinische Biophysik, Urologische Klinik und Poliklinik, Technische Universität, Ismaningerstr. 22, D-8000 München 8, Germany
 SO Acta Pharmaceutica, (1992) 42/4 (263-271).
 ISSN: 1330-0075 CODEN: ACPHEE
 CY Croatia
 DT Journal; Conference Article
 FS 029 Clinical Biochemistry
 LA English
 SL English; Slovenian
 AB **Lipid** membranes are being used increasingly in many biotechnological and biological systems. For the success of such applications it is crucial to understand and quantify the interactions between **lipid** bilayers and soluble macromolecules, however. We argue that the efficiency of specific binding of any large molecule (e.g. an antibody) to its ligand attached to a **lipid** membrane is diminished by the proximity of the bilayer surface. This appears to be due to the same non-Coulombic, hydration dependent short-range force which also prevents the colloidal collapse of a stack of **lipid** membranes. The strength of this force, as a rule, decreases with decreasing polarity of the **lipid** headgroups. Nonspecific protein adsorption of the **lipid** bilayers, on the contrary, is little affected by the overall repulsion between a macromolecule and the bilayer surface; rather, this adsorption is governed by the number of defects and/or by the available hydrophobic binding sites in a given membrane. Multiple nonspecific protein adsorption, which results in the macromolecular denaturation, is one of the main reasons for the rapid elimination of **lipid** vesicles from the blood stream in vivo. To promote the longevity of liposomes it is therefore necessary to modify the surfaces of their constituent **lipid** bilayers. Increasing the surface density of the net charges and/or the bilayer surface hydrophilicity is of little use in this respect; to affect the nonspecific bilayer-protein interactions significantly, an optimal number of water soluble, short **polymers** should be attached to the **lipid** headgroups. These increase the repulsive barrier of the **lipid** membrane surface and often may prolong the life-time of the resulting **lipid** vesicles dramatically. Vesicles consisting of a 8/2 (mol mol-1) mixture of distearoylphosphatidylethanolamine-PEG (**DSPE** -PEG) with distearoylphosphatidylcholine or of a 7/3 (mol mol-1) mixture **DSPE**-PEG and soy-bean phosphatidylcholine, for example, remain in the murine circulation 8 to 10 times longer than standard phosphatidylcholine liposomes. The **polymer**-coated long-lived liposomes, moreover, have a lower affinity for protein binding, induce less protein denaturation, are not very attractive to phagocytic cells, and also accumulate highly efficiently in tumors.

L43 ANSWER 23 OF 27 MEDLINE
 AN 95178599 MEDLINE
 DN 95178599 PubMed ID: 7873652
 TI New amphipatic **polymer-lipid** conjugates forming
 long-circulating reticuloendothelial system-evading liposomes.
 AU Woodle M C; Engbers C M; Zalipsky S
 CS Liposome Technology, Inc., Menlo Park, California 94025.
 SO BIOCONJUGATE CHEMISTRY, (1994 Nov-Dec) 5 (6) 493-6.
 Journal code: 9010319. ISSN: 1043-1802.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199504
 ED Entered STN: 19950419
 Last Updated on STN: 19950419
 Entered Medline: 19950404
 AB **Lipid**-conjugates of two amphipatic **polymers**,
 poly(2-methyl-2-oxazoline) (PMOZ) and poly(2-ethyl-2-oxazoline) (PEOZ)
 (degree of **polymerization** approximately 50) were synthesized by
 linking glutarate esters of the **polymers** to
 distearoylphosphatidylethanolamine (**DSPE**) or alternatively by
 termination of the **polymerization** process with **DSPE**.
 Surface-modified liposomes (90 +/- 5 nm) prepared from either conjugate (5
 mol % of total **lipid**) were injected into rats and followed by
 blood level and tissue distribution measurements. Both **polymers**
 PEOZ and PMOZ were found to convey long circulation and low hepatosplenic
 uptake to liposomes to the same extent as polyethylene glycol (PEG), the
 best known material for this purpose. This is the first demonstration of
 protection from rapid recognition and clearance conveyed by alternative
polymers, which is equal to the effect of PEG.

L45 ANSWER 16 OF 23 BIOTECHDS COPYRIGHT 2003 THOMSON DERWENT AND ISI
 AN 1999-00209 BIOTECHDS
 TI Processes for drug delivery, therapeutic imaging, and diagnostic imaging;
 used for targeted drug delivery and gene therapy
 AU Unger E C
 PA Imarx-Pharm.
 LO Tucson, AZ, USA.
 PI WO 9842383 1 Oct 1998
 AI WO 1998-US3632 25 Feb 1998
 PRAI US 1997-925353 8 Sep 1997; US 1997-823791 21 Mar 1997
 DT Patent
 LA English
 OS WPI: 1998-557028 [47]
 AB A method for delivering a bioactive agent (BA) to a patient is claimed,
 involving administration of a composition containing an **anionic**
lipid, a cationic counter ion, a **lipid** covalently
 bonded to a **polymer**, and the BA. Therapeutic ultrasound is
 then applied to facilitate delivery of the BA in a desired region. An
 alternate method involves administering a compound containing a charged
lipid in place of the **anionic lipid**, a
 counter ion in place of the cationic counter ion, and an additional
 targeting ligand. Also claimed is a means of providing an image of an
 internal region of a patient by administering the alternate compound and
 scanning the patient using diagnostic imaging to produce visible images
 of the desired region. The claims also cover a contrasting agent,
 composed of the constituents of the alternate composition. These
 compositions and processes are for use in the targeted delivery of drugs,
 with improved efficacy and reduced toxicity. The preferred BA is a
 diagnostic agent, or genetic material, e.g. recombinant **nucleic**
 acid, antisense **nucleic** acid, hammerhead RNA, ribozyme,
 hammerhead ribozyme, oligonucleotide, etc. (167pp)

L48 ANSWER 9 OF 11 MEDLINE DUPLICATE 2
 AN 2001206621 MEDLINE
 DN 21144495 PubMed ID: 11249656
 TI Recent developments in mucosal delivery of pDNA vaccines.
 AU Barnes A G; Barnfield C; Brew R; Klavinskis L S
 CS Department of Immunobiology, Guy's, King's and St Thomas' Medical School
 (GKT), St Thomas Street, London SE1 9RT, UK.. linda.klavinskis@kcl.ac.uk
 SO Curr Opin Mol Ther, (2000 Feb) 2 (1) 87-93. Ref: 60
 Journal code: 100891485. ISSN: 1464-8431.
 CY England: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 General Review; (REVIEW)
 (REVIEW, TUTORIAL)
 LA English
 FS Priority Journals
 EM 200104
 ED Entered STN: 20010417
 Last Updated on STN: 20010417
 Entered Medline: 20010412
 AB The development of **DNA** vaccination to mucosal surfaces has
 continued apace over the last 2 years, with the investigation of several
 novel delivery vehicles. There have been advances in the understanding of
 the basic immunological mechanisms behind the induction of immune
 responses by **plasmid DNA**. The mechanistic insights are
 paving the way for the design of a second generation of mucosally
 delivered **DNA** vaccines. This article reviews the recent progress
 in the field of **microparticle**, cationic **lipid** and
 bacterial delivery systems. All these mechanisms afford some protection
 from environmental degradation and facilitate **DNA** uptake. These
 methods have been compared with respect to transfection efficiency,
 ability to elicit a full range of immune responses and their relative
 safety for in vivo applications.

L48 ANSWER 6 OF 11 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 2001:439332 BIOSIS
DN PREV200100439332
TI Methods of forming protein-linked lipidic microparticles, and compositions thereof.
AU Papahadjopoulos, Demetrios; Hong, Keelung; Zheng, Weiwen; Kirpotin, Dmitri B. (1)
CS (1) San Francisco, CA USA
ASSIGNEE: The Regents of the University of California
PI US 6210707 April 03, 2001
SO Official Gazette of the United States Patent and Trademark Office Patents, (Apr. 3, 2001) Vol. 1245, No. 1, pp. No Pagination. e-file.
ISSN: 0098-1133.
DT Patent
LA English
AB The present invention provides for **lipid:nucleic acid** complexes that have increased shelf life and high transfection activity in vivo following intravenous injection, and methods of preparing such complexes. The methods generally involve contacting a **nucleic acid** with an organic polycation to produce a condensed **nucleic acid**, and then combining the condensed **nucleic acid** with a **lipid** comprising an amphiphilic cationic **lipid** to produce the **lipid:nucleic acid** complex. This complex can be further stabilized by the addition of a hydrophilic **polymer** attached to hydrophobic side chains. The complex can also be made specific for specific cells, by incorporating a targeting moiety such as an Fab' fragment attached to a hydrophilic **polymer**. The present invention further relates to lipidic microparticles with attached proteins which have been first conjugated to linker molecules having a hydrophilic **polymer** domain and a hydrophobic domain capable of stable association with the **microparticle**, or proteins which have been engineered to contain a hydrophilic domain and a **lipid** moiety permitting stable association with the **microparticle**.